

## DESCRIPTION

MONOCLONAL ANTIBODY INDUCING APOPTOSIS

## TECHNICAL FIELD

5 This invention relates to novel monoclonal antibodies having the property of inducing apoptosis of nucleated blood cells with Integrin Associated Protein (IAP), as well as to ~~their fragments, peptides and low~~  
10 ~~molecular compounds,~~ and to hybridomas that produce the monoclonal antibodies. The novel antibodies are useful as therapeutic agents for myeloid leukemia and lymphoid leukemia.

## BACKGROUND ART

Granulocyte colony-stimulating factors, such as recombinant granulocyte colony-stimulating factor (rG-CSF), have been known in the prior art as humoral factors that stimulate differentiation and proliferation of granulocytes. Reports based on *in vivo* experiments with mice have shown that administration of rG-CSF results in not only  
20 accelerated myelopoiesis in bone marrow but also notable extramedullary hemopoiesis in the spleen, and proliferation of all hemopoietic precursor cells, including hemopoietic stem cells, in the spleen. The mechanism of such extramedullary hemopoiesis in the spleen has been believed  
25 that stimulation by rG-CSF alters the hemopoietic microenvironment of the spleen and promotes the hemopoiesis supporting ability thereof, thus inducing hemopoiesis.

In order to elucidate the hemopoietic function in the spleen, the present inventors have previously focused on stromal cells of the spleen following repeated administration of rG-CSF. The inventors have made efforts to examine how the hemopoietic function is promoted by rG-CSF via stromal cells, and have established a hemopoietic stromal cell line (CF-1 cells) from mouse spleen by repeated administration of rG-CSF. The inventors have studied the hemopoiesis-supporting ability of the hemopoietic stromal cells and confirmed the colony-stimulating activity *in vitro* and the hemopoietic stem cell-supporting ability *in vivo* [Blood, 80, 1914 (1992)].

However, while one cell line of the splenic stromal cells has been established (CF-1 cells) and its cytological characteristics have been studied, specific antibodies that recognize the surface antigens of these cells have never been prepared, nor have their characteristics been elucidated yet in any way.

#### DISCLOSURE OF INVENTION

In light of the aforementioned findings relating to splenic stromal cells and the results of prior research, the present inventors have earnestly made further research aiming at developing specific antibodies that can recognize the splenic stromal cells, made efforts to prepare monoclonal antibodies using the aforementioned splenic stromal cell line as a sensitizing antigen, and finally succeeded in obtaining novel monoclonal antibodies.

The inventors have further studied the properties of the monoclonal antibodies obtained as above and found that the monoclonal antibodies have the property of inducing myeloid cell apoptosis. These monoclonal antibodies have been designated "BMAP-1 antibody", which will be hereinafter referred to as such.

The inventors have also examined the antigen recognized by BMAP-1 antibody and found that it is mouse Integrin Associated Protein (mouse IAP) (GenBank, Accession Number Z25524) by direct expression cloning.

The action of BMAP-1 antibodies has been studied using recombinant cells into which the gene for mouse IAP had been introduced. Specifically, the mouse IAP gene was introduced into mouse Jurkat cells, which did not express mouse IAP, by a conventional method to create a mouse IAP-expressing cell line (recombinant Jurkat cells), and the action of BMAP-1 antibody on the mouse IAP-expressing cells has been investigated by MTS assay and DNA fragmentation by using flow cytometry (Japanese Patent Application No. HEI 9-67499).

It has been expected upon these findings that monoclonal antibodies for the antigen of human Integrin Associated Protein (hereinafter referred to as human IAP; amino acid sequence and base sequence described in J. Cell Biol., 123, 485-496, 1993; see also Journal of Cell Science, 108, 3419-3425, 1995) should have an effect of inducing apoptosis of nucleated blood cells that express this antigen

(myeloid cells and lymphocytes), and the present inventors have made efforts to prepare monoclonal antibodies for the antigen of human Integrin Associated Protein and succeeded in obtaining monoclonal antibodies that induce apoptosis of human nucleated blood cells expressing this antigen.

In other words, it is an object of this invention to provide novel monoclonal antibodies having the property of inducing apoptosis of nucleated blood cells (myeloid cells and lymphocytes) with human Integrin Associated Protein (human IAP), and fragments thereof, as well as hybridomas that produce the monoclonal antibodies.

These novel monoclonal antibodies are useful as therapeutic agents for myeloid leukemia and lymphoid leukemia.

The reported functions of Integrin Associated Protein are the action of binding with the  $\beta$  chain of integrin  $\alpha V \beta 3$  to support binding between  $\alpha V \beta 3$  and its ligand vitronectin (J. Cell. Biol., 123, 485-496 (1993)), that of inducing inflow of  $Ca^{2+}$  into the vascular endothelium upon adhesion of neutrophils with the vascular endothelium (J. Biol. Chem., 268, 19931-19934 (1993)), and that of supporting migration of neutrophils through the vascular endothelium (Proc. Natl. Acad. Sci. USA, 92, 3978-3982 (1995)), but no reports have been published on its function relating to apoptosis of nucleated blood cells.

The monoclonal antibodies of the invention are antibodies that specifically recognize human Integrin

Associated Protein. They therefore exhibit a function of distinguishing and identifying human Integrin Associated Protein.

5 In addition, the monoclonal antibodies of the invention are antibodies that exhibit the property of inducing apoptosis of nucleated blood cells (myeloid cells and lymphocytes) with human Integrin Associated Protein. Apoptosis is a phenomenon in which nuclear chromatin DNA is cleaved into nucleosome units (known as a "ladder formation"), resulting in death of the cell and which is also referred to as cell suicide.

Monoclonal antibodies hitherto known to have the property of inducing apoptosis of nucleated blood cells (myeloid cells and lymphocytes) include anti-Fas antibody (Cell, 66; 233-243, 1991), anti-CD43 antibody (Blood, 86, 502-511, 1995) and anti-HLA Class Ia1 Domain antibody (Blood, 90, 726-735, 1997), but the property of inducing apoptosis of nucleated blood cells by the Integrin Associated Protein-recognizing antibodies of this invention  
20 has never been known. The monoclonal antibodies of the invention are therefore defined as encompassing any monoclonal antibody capable of specifically recognizing Integrin Associated Protein and having the property of inducing apoptosis of nucleated blood cells (myeloid cells  
25 and lymphocytes) with Integrin Associated Protein.

The antibodies of the invention are not limited only to those that induce apoptosis of all nucleated blood

cells. They also include those that induce apoptosis of at least one type of nucleated blood cells. Specifically, it is sufficient in the case of myeloid leukemia to induce apoptosis of at least myeloid cells.

5 More specifically, this invention provides monoclonal antibodies that induce apoptosis of nucleated blood cells having Integrin Associated Protein (IAP).

10 <sup>A</sup> The invention further provides ~~fragments, peptides~~  
~~and low molecular compounds of monoclonal antibodies~~ that induce apoptosis of nucleated blood cells having Integrin Associated Protein (IAP).

15 The invention still further provides hybridomas that produce the monoclonal antibodies.

20 The invention still further provides an antileukemic agent that contains a substance that binds to IAP and promotes the action of IAP to induce apoptosis of nucleated blood cells.

25 The invention still further provides an antileukemic agent characterized in that the substance is a monoclonal antibody.

<sup>A</sup> <sup>A3</sup> The invention still further provides an antileukemic agent characterized in that the substance is a ~~fragment, a peptide or a low molecular compound of the~~  
~~monoclonal antibodies.~~

#### 25 BRIEF DESCRIPTION OF DRAWINGS

Fig. 1 is an electrophoresis pattern showing a band for human IAP amplified by PCR using cDNA prepared from

mRNA of HL-60 cell line. From left are shown a molecular weight marker (M), human IAP (1) and  $\beta$ -actin (2).

Fig. 2 is a graph showing the level of expression of human IAP by L1210 cells that have expressed human IAP, using anti-CD47 antibody. The peak represents L1210 cells transfected with only pCOS1 gene as a control.

Fig. 3 is another graph showing the level of expression of human IAP by L1210 cells that have expressed human IAP, using anti-CD47 antibody. The peak shows that human IAP expression has definitely increased in L1210 cells transfected with the human IAP gene.

Fig. 4 is a graph showing antibody titers in immunized mice. The left peak represents intact L1210 cells. The right peak represents L1210 cells transfected with human IAP, showing that the serum of the mouse subjected to cell fusion clearly recognizes human IAP.

Fig. 5 is a bar graph showing the results of a growth inhibition experiment (Jurkat cells) using a hybridoma culture supernatant.

Fig. 6 is a bar graph showing the results of a growth inhibition experiment (ARH77 cells) using a hybridoma culture supernatant.

Fig. 7 is a graph showing the apoptosis-inducing effect on Jurkat cells by a culture supernatant (as analyzed by PI staining), which is the result for an 8G2 culture supernatant used as a control. R1 indicates the percentage (%) of apoptosis, which is 7.43%.

Fig. 8 is a graph showing the apoptosis-inducing effect on Jurkat cells by a culture supernatant (as analyzed by PI staining), which is the result for 7D2-E3. R1 indicates the percentage (%) of apoptosis, which is 9.84%.

5 Fig. 9 is a graph showing the apoptosis-inducing effect on Jurkat cells by a culture supernatant (as analyzed by PI staining), which is the result for 11C8. R1 indicates the percentage (%) of apoptosis, which is 15.32%.

10 Fig. 10 is a graph showing the apoptosis-inducing effect on HL-60 cells by a culture supernatant (as analyzed by PI staining), which is the result for an 8G2 culture supernatant used as a control. M1 indicates the percentage (%) of apoptosis, which is 6.94%.

15 Fig. 11 is a graph showing the apoptosis-inducing effect on HL-60 cells by a culture supernatant (as analyzed by PI staining), which is the result for 11C8. M1 indicates the percentage (%) of apoptosis, which is 12.16%.

20 Fig. 12A is a monochrome photomicrograph showing the result of apoptosis analysis (TUNEL method) in a coculturing system with KM-102 and HL-60 cells, using 9C5 culture supernatant as a control. The apoptotic cells are stained black or brown. The nuclear staining was accomplished with Methyl Green, and the magnification is 100x.

25 Fig. 12B is a color photomicrograph showing the result of apoptosis analysis (TUNEL method) in a coculturing system with KM-102 and HL-60 cells, using 9C5 culture



supernatant as a control. The apoptotic cells are stained black or brown. The nuclear staining was accomplished with Methyl Green, and the magnification is 100x.

Fig. 13A is a monochrome photomicrograph showing the result of apoptosis analysis (TUNEL method) in a coculturing system with KM-102 and HL-60 cells, using 11C8 culture supernatant. More TUNEL-positive cells are seen than in Fig. 12. The apoptotic cells are stained black or brown. The nuclear staining was accomplished with Methyl Green, and the magnification is 100x.

Fig. 13B is a color photomicrograph showing the result of apoptosis analysis (TUNEL method) in a coculturing system with KM-102 and HL-60 cells, using 11C8 culture supernatant. More TUNEL-positive cells are seen than in Fig. 12. The apoptotic cells are stained black or brown. The nuclear staining was accomplished with Methyl Green, and the magnification is 100x.

Fig. 14 is an electrophoresis pattern showing the results of SDS-PAGE analysis of IgG purified from hybridoma lines 7D2-E3 and 11C8. Shown are molecular weight markers (M, M'), mouse IgG (authentic sample) under non-reducing conditions (1), 7D2-E3 (2), 11C8 (3), mouse IgG (authentic sample) under reducing conditions (4), 7D2-E3 (5) and 11C8 (6).

Fig. 15 shows the results of analysis of CD47 expression by flow cytometry, using HL-60 cells.

Fig. 16 shows the results of analysis of CD47 expression by flow cytometry, using Jurkat cells.

Fig. 17 shows results for mIgG (10 µg/ml) as a control to demonstrate its apoptosis-inducing effect on L1210 cells transfected with the human IAP gene (L1210-hIAP) (incubation for 72 hours).

Fig. 18 shows the apoptosis-inducing effect of MABL-1 (10 µg/ml) on L1210 cells transfected with the human IAP gene (incubation for 72 hours).

Fig. 19 shows the apoptosis-inducing effect of MABL-2 (10 µg/ml) on L1210 cells transfected with the human IAP gene (incubation for 72 hours).

Fig. 20 shows results for mIgG (10 µg/ml) as a control to demonstrate its apoptosis-inducing effect on Jurkat cells (incubation for 48 hours).

Fig. 21 shows the apoptosis-inducing effect of MABL-1 (10 µg/ml) on Jurkat cells (incubation for 48 hours).

Fig. 22 shows the apoptosis-inducing effect of MABL-2 (10 µg/ml) on Jurkat cells (incubation for 48 hours).

Fig. 23 shows results for mIgG (10 µg/ml) as a control to demonstrate its apoptosis-inducing effect on L1210 cells transfected with the human IAP gene introduced therein (L1210-hIAP) (incubation for 72 hours).

Fig. 24 shows the apoptosis-inducing effect of MABL-2 Fab fragments (10 µg/ml) on L1210 cells transfected with the human IAP gene.

Fig. 25 is an SDS electrophoresis pattern for MABL-2 antibody Fab fragments.

Fig. 26 shows a notably extended survival period upon treatment with MABL-2.

Fig. 27 shows the results of ELISA for Example 5(2).

Fig. 28 shows a notably extended survival period upon treatment with MABL-2 F(ab')<sub>2</sub> fragments.

Fig. 29 is an SDS electrophoresis pattern for MABL-1 antibody and MABL-2 antibody F(ab')<sub>2</sub> fragments.

Fig. 30 shows that human IgG levels of mouse serum were decreased significantly in the groups treated with MABL-1 and MABL-2, which indicates anti-tumor effects of these antibodies.

#### **BEST MODE FOR CARRYING OUT THE INVENTION**

##### Preparation of Monoclonal Antibody

The monoclonal antibodies of this invention can generally be prepared in the following manner. That is, monoclonal antibodies of the invention may be obtained, for example, by using human Integrin Associated Protein as the sensitizing antigen, immunizing animals with the antigen by an immunization method known in the art, performing cell fusion by a cell fusion method known in the art and cloning by a cloning method known in the art.

More specifically, a preferable method of preparing monoclonal antibodies of the invention is, for example, a method wherein recombinant cells of the mouse

leukemia cell line L1210 that express human Integrin Associated Protein are used as the sensitizing antigen, plasma cells (immunocytes) of a mammal immunized with the sensitizing antigen are fused with myeloma cells of mammals such as mice, the resulting fused cells (hybridomas) are cloned, the clones producing the antibodies of the invention that recognize the aforementioned cell line are selected from the resulting clones and cultured, and the target antibodies are obtained.

The above method is merely one possible example of the invention and, for example, the sensitizing antigen is not limited to the aforementioned L1210 recombinant cells but may also be human Integrin Associated Protein (IAP) itself, or human IAP in soluble form; the target monoclonal antibodies that induce apoptosis of nucleated blood cells (myeloid cells and lymphocytes) can be prepared in the same manner as in the L1210 recombinant cells mentioned above.

The phage display method may also be used to prepare a target monoclonal antibody from a cDNA library for the antibody.

The mammals to be immunized with the sensitizing antigen in the method of preparing the monoclonal antibodies are not particularly limited, but they are preferably selected in consideration of their compatibility with the myeloma cells used for cell fusion, and mice, rats, hamsters and the like are general suitable.

The immunization is preferably accomplished by a standard method. For example, the human Integrin Associated Protein-expressing L1210 recombinant cells are administered to the animal by intraperitoneal injection or the like.

5 More specifically, an appropriate dilution or suspension with PBS or physiological saline is preferably administered to the animal a few times at 10-day intervals. The immunocytes used are preferably spleen cells extracted after the final administration of the cells.

10 The mammalian myeloma cells used as the parent cells for fusion with the immunocytes may be any of various cell lines known in the art, for example, P3 (P3X63Ag8.653) [J. Immunol., 123, 1548 (1978)], P3-U1 [Current Topics in Microbiology and Immunology, 81, 1-7 (1978)], NS-1 [Eur. J. Immunol., 6, 511-519 (1976)], MPC-11 [Cell, 8, 405-415 (1976)], Sp2/0-Ag14 [Nature, 276, 269-270 (1978)], FO [J. Immunol. Meth., 35, 1-21 (1980)], S194 [J. Exp. Med., 148, 313-323 (1978)] and R210 [Nature, 277, 131-133 (1979)].

20 The cell fusion between the immunocytes and myeloma cells may be performed basically according to a conventional method, such as the method of Milstein et al. [Methods Enzymol., 73, 3-46 (1981)].

25 More specifically, the cell fusion is carried out, for example, in a common nutrient medium in the presence of a fusion promoter. For example, the fusion promoter used may be polyethylene glycol (PEG), Sendai virus (HVJ) or the like, and, if desired, an adjuvant such as dimethyl

sulfoxide may also be added appropriately in order to increase fusion efficiency. The immunocytes are used preferably in the amount of 1-10 times as much as myeloma cells. The medium used for the cell fusion may be, for example, RPMI-1640 medium, MEM medium and the like, which are suitable for growth of myeloma cell lines, or other media commonly used for such cell culturing, and it may also be used in combination with a serum supplement such as fetal bovine serum (FBS).

The cell fusion is carried out by thoroughly mixing prescribed amounts of the immunocytes and myeloma cells in the medium, adding a solution of PEG preheated to about 37°C, the PEG having an average molecular weight of approximately 1,000-6,000, for example, to the medium usually at a concentration of about 30-60% (W/V), and mixing. A suitable medium is then successively added, and the supernatant obtained by centrifugation is removed. This procedure is repeated to produce the target hybridomas.

The hybridomas are selected by culturing in a common selection medium, such as HAT medium (a medium containing hypoxanthine, aminopterin and thymidine). Culturing in the HAT medium is continued for a sufficient time to allow death of all the cells other than the target hybridomas (all the non-fused cells), which is usually from a few days to a few weeks. The usual limiting dilution method is then employed for screening and monocloning of hybridomas producing the target antibodies.

The hybridomas prepared in this manner that produce the monoclonal antibodies of the invention may be subcultured in common medium, and may be placed in long-term storage in liquid nitrogen.

5 In order to obtain the monoclonal antibodies of the invention from the hybridomas, any suitable methods may be employed, such as a method wherein the hybridomas may be cultured according to standard methods and the antibodies may be obtained from the culture supernatants; or alternatively, a method wherein the hybridomas may be administered to a compatible mammal for proliferation and then the antibodies may be obtained from the ascites fluid thereof. The former method is suitable for obtaining highly pure antibodies, while the latter method is more suited for mass production of antibodies.

The antibodies obtained by the aforementioned methods can be highly purified by utilizing standard purification methods such as salting-out, gel filtration, affinity chromatography, or the like.

#### 20 Monoclonal Antibody Fragments

25 The monoclonal antibodies of this invention may be the complete antibodies described above, or fragments thereof. That is, they may be any fragments of a monoclonal antibody of the invention that specifically recognize human Integrin Associated Protein and induce apoptosis of nucleated blood cells (myeloid cells and lymphocytes) having human Integrin Associated Protein. Such fragments include

Fab, F(ab')<sub>2</sub>, Fab', etc. These fragments can be prepared by digestion with an enzyme such as papain, pepsin, ficin or the like. The properties of the obtained fragments can be confirmed in the same manner as described above.

5 Peptides and Low Molecular <sup>weight</sup> Compounds Having the Same Function as the Monoclonal Antibodies

The monoclonal antibodies described above, which recognize human Integrin Associated Protein and induce apoptosis of nucleated blood cells, also encompass peptides and low molecular <sup>weight</sup> compounds that likewise recognize IAP and induce apoptosis of nucleated blood cells.

Properties of Monoclonal Antibodies of the Invention

As specifically described in the following Examples, the monoclonal antibodies of the invention specifically recognize human Integrin Associated Protein.

The monoclonal antibodies of the invention also induce apoptosis of nucleated blood cells (myeloid cells and lymphocytes) with human Integrin Associated Protein.

20 These properties can be utilized to obtain useful therapeutic agents in the field of treatment for myeloid leukemia and lymphoid leukemia.

25 Thus, it will be readily appreciated that the construction of specific systems involving the use of the monoclonal antibodies of the invention, as antibodies to specifically recognize an antigen that causes apoptosis of nucleated blood cells, for distinction and identification of the antigens, or the use of the unique properties of the



monoclonal antibodies as therapeutic agents for myeloid leukemia and lymphoid leukemia, as well as any modifications and applications of the system, are also within the scope of this invention insofar as they can be carried out by applying standard methods that are obvious to those skilled in the art.

#### Antileukemic Agents

An antileukemic agent according to this invention is based on the fact that the action of IAP is promoted by binding of an antibody or the like of the invention. While there are no particular limitations on the dose of the antibody of the invention, it is preferably in the range of 5  $\mu$ g to 500 mg/kg.

#### **EXAMPLES**

This invention will now be explained in greater detail by way of the following examples; however, the invention is not to be limited to these examples.

##### Example 1 (Monoclonal Antibody Preparation)

##### (1) Sensitizing antigen and immunization method

Antigen sensitization was accomplished using a recombinant cell line as the sensitizing antigen, which was the L1210 cells transfected with human IAP gene and highly expressed the product. L1210 is obtained from the DBA mouse-derived leukemia cell line (ATCC No. CCL-219, J. Natl. Cancer Inst. 10:179-192, 1949).

The human IAP gene was amplified by PCR using a primer with a human IAP-specific sequence (sense primer:

GCAAGCTTATGTGGCCCCCTGGTAGCG, antisense primer:

GCGGCCGCTCAGTTATTCCTAGGAGG) and cDNA prepared from mRNA of HL-60 cell line (Clontech laboratories, Inc.) as the template (Fig. 1).

5           The PCR product was subcloned into a cloning vector pGEM-T (Promega Corporation) and used to transform *E. coli* JM109 (Takara Shuzo Co., Ltd.), and after confirming the nucleotide sequence of the insert DNA with a DNA sequencer (373A DNA Sequencer, available from ABI), it was subcloned with an expression vector pCOS1.

10           Expression vector pCOS1 is a derivative of pEF-BOS (Nucleic Acids Research, 18, 5322, 1990), and it is a vector obtained by subcloning the neomycin resistant gene using human elongation factor-1 $\alpha$  as a promoter/enhancer. This human IAP-subcloned expression vector was used for gene introduction into L1210 cell line with DMRIE-C (GIBCO/BRL), selection was performed with Geneticin (final concentration: 1 mg/ml, available from GIBCO/BRL), and the gene-introduced L1210 cells were cloned by the limiting dilution method.

20           The antigen expression of the obtained clones was examined using human IAP-recognizing anti-CD47 antibody (PharMingen), and the clones with high levels of expression were selected as antigen-sensitized cells (Figs. 2, 3). For culturing of the recombinant L1210 cells, 10% fetal bovine serum (FBS, available from Moregate Inc.) and Iscove's-  
25           Modified Dulbecco's Medium (IMDM) (GIBCO/BRL) were used as

the medium, and the cells were subcultured in a 5% CO<sub>2</sub> incubator at a temperature of 37°C.

The immunized animals used were DBA/2 mice (bred by Charles River, Japan), which were of the same strain as the L1210 cells. The human Integrin Associated Protein (IAP) gene-transfected L1210 cells, used for antigen sensitization, were incubated for about 30 min with mitomycin C (Kyowa Hakko Kogyo Co., Ltd.) at a concentration of 200 µg/ml, and after suspending growth of the cells, mitomycin C was thoroughly washed off prior to suspension in PBS.

The cells were intraperitoneally injected into the mice three times at intervals of about 10 days, at approximately  $5 \times 10^6$  cells each time. After the third immunization, blood was taken from the eye socket, the serum was diluted 50-fold with PBS containing 1% BSA, and binding between the diluted serum and the recombinant L1210 cells used for antigen sensitization was confirmed with a FACScan (Becton Dickinson and Company) (Fig. 4); the mouse having the best antiserum activity was subjected to a booster immunization with intraperitoneal injection of  $1 \times 10^7$  cells 5 days after the fourth immunization. Four days after the final immunization, the mouse was sacrificed and the spleen extracted.

## (2) Cell fusion

After thinly slicing the spleen extracted from the mouse, the dissociated spleen cells were centrifuged and

then suspended in IMDM medium, allowed to float, and thoroughly rinsed. Separately, the mouse myeloma cell line P3-U1 [Current Topics in Microbiology and Immunology, 81, 1-7 (1978)] was cultured in IMDM medium containing 10% fetal bovine serum (FBS, available from Moredgate Inc.), and after rinsing similarly with the IMDM medium, the  $1 \times 10^7$  cells were placed in a centrifuge tube in admixture with  $5 \times 10^7$  cells of the spleen cells and subjected to cell fusion according to a standard method [Clin. Exp. Immunol., 42, 458-462 (1980)], using polyethylene glycol 4000 (Nakarai Chemical Co., Ltd.).

The resulting fused cells were then suspended in IMDM medium containing 10% FBS and a fused cell growth stimulating agent (BM-Condimed H1, available from Boehringer Mannheim Biochemicals) and dispensed into a 96-well plate for culturing at 37°C in a 5% CO<sub>2</sub> incubator. On the following day, the cells were placed in the HAT selection medium and then the 10% FBS/IMDM medium containing the growth-stimulating agent, and culturing was continued to sustain growth.

In order to examine the effect of the culture supernatant of these fused cells on leukemia cell lines, the medium for fused cells was replaced with IMDM medium containing 10% FBS, and culturing was continued to sustain growth.

(3) Screening

The following screening was performed using the culture supernatant of the aforementioned fused cells.

[1] Primary screening

Cells of a mouse spleen stromal cell line (CF-1 cells) transfected with the human Integrin Associated Protein (IAP) gene (recombinant cells into which the same plasmid was subcloned as the plasmid used to prepare the human IAP-expressing L1210 cells used for antigen sensitization) were seeded in a 96-well plate at  $1 \times 10^4$  cells per well and cultured overnight, and then fixed with 2% PLP (periodate-lysine-paraformaldehyde) to prepare an ELISA plate. After rinsing, the plate was subjected to blocking for 1 h at room temperature using a 1% BSA solution, and after further rinsing, 50  $\mu$ l of the culture supernatant of each hybridoma was added for incubation at room temperature for one hour.

After rinsing, anti-mouse IgG+A+M (H+L) (Zymed Laboratories Inc.) labeled with alkaline phosphatase was added prior to incubation at room temperature for 1 h.

After rinsing, SIGMA 104 substrate (Sigma-Aldrich Corporation) was added to provide a final concentration of 1 mg/ml, incubation was continued at room temperature, and the specific activity was measured with a microplate reader (Model 3550, available from BioRad Laboratories Inc.).

As a result, appearance of hybridomas was confirmed in 2089 wells among the hybridomas seeded in 2880 wells, with 187 wells being positive in the primary

screening. 50  $\mu$ l each of Mouse IgG1 as a negative control and anti-human CD47 antibody (BD PharMingen) as a positive control were added at a concentration of 3  $\mu$ g/ml, respectively, prior to incubation at room temperature for 1 h.

## [2] Secondary screening

The clones judged as positive in the primary screening were subjected to an ELISA system using human Integrin Associated Protein (IAP)-expressing CF-1 cells, where the negative control was CF-1 cells transfected with only the expression vector pCOS1, in order to screen whether the antibodies produced by the hybridomas would specifically recognize human IAP.

As a result, the positive was confirmed for 21 of the 187 wells found to be positive in the primary screening. Table 1 shows the specific binding of human IAP with 7D2 and 11C8 as representative examples among these, in terms of the absorbance in ELISA.

(Table 1) ELISA analysis of specific binding of hybridoma culture supernatants with human IAP

Table 1

<Raw data>	PBS	3 µg/ml		
		αhCD47	7D2	11C8
CF1-pCOS1	0.185	0.160	0.189	0.149
CF1-hIAP-55-8	0.192	0.456	0.568	0.812
<Subtracted>	PBS	3 µg/ml		
		αhCD47	7D2	11C8
Specific binding	0.007	0.296	0.379	0.663

### [3] Tertiary screening

The clones judged to be positive in the secondary screening were subjected to a growth inhibition test using Jurkat cells (human T cell lymphoma line) and ARH77 cells (human myeloma cell line). 100 µl of the Jurkat cells at  $5 \times 10^3$  cells per well and the ARH77 cells at  $1 \times 10^4$  cells per well were seeded in each well of a 96-well plate, and 5 or 10 µl of culture supernatant of the hybridoma clones were added to the cell suspensions. After culturing for about 2 days, the cell numbers were measured by MTS assay. As a control, 5 or 10 µl each of IMDM medium containing 10% FBS and culture supernatants of clones that were negative in the primary screening (8G2 and 9C5) were added.

Figs. 5 and 6 show the results of the growth inhibition effect of four representative clones, 11C8, 7D2-E3 (subclone of 7D2), 13F1 and 2F12.

#### (4) Antibody properties

[1] The immunoglobulin types of the culture supernatants of 11C8, 7D2-E3, 13F1 and 2F12 were examined using an ELISA system.

Specifically, human Integrin Associated Protein (IAP)-expressing CF-1 cells were seeded into a 96-well plate to prepare an ELISA plate, and then 50  $\mu$ l of each culture supernatant was added, alkaline phosphatase-labeled anti-mouse IgG antibody (Zymed Laboratories Inc.) or anti-mouse IgM antibody (Biosource Intl., Inc.) were reacted therewith as secondary antibodies, and the activity was measured with a microplate reader. As a result, 11C8 and 7D2-E3 were confirmed to be IgG, while 13F1 and 2F12 were confirmed to be IgM.

[2] The DNA fragmentation of the two clones 11C8 and 7D2-E3 among the four clones described above was analyzed by flow cytometry (FACScan, available from Becton, Dickinson and Company) using Jurkat cells and HL-60 cells. The Jurkat cells were used for 11C8 and 7D2-E3, and the HL-60 cells were used for 11C8.

The Jurkat cells and HL-60 cells were seeded in a 12-well plate at  $4 \times 10^4$  cells per well/2 ml, respectively, and 200  $\mu$ l of the culture supernatants of 7D2-E3 and 11C8 were added. The cells were cultured for 2 days, and



measured. As a control, 8G2 culture supernatant was added in an equal volume. The cells were recovered from the culturing plate and a cell pellet was fixed under 200 x g for 60 minutes at 4°C in 2 ml of chilled 70% ethanol. The cells were then centrifuged, rinsed in 1 ml of PBS and resuspended in 0.5 ml of PBS. To a 0.5 ml sample of the cells, 0.5 ml of RNase (Type I-A, Sigma-Aldrich Corporation, St. Louis, MO, USA; 1 mg/ml in PBS) was added, and these were mixed with a 1 ml propidium iodide solution (PI, Sigma, 100 µg/ml in PBS). The mixed cells were incubated for 60 min in a darkroom at 37°C, and then kept in the darkroom at 4°C and measured by flow cytometry.

As shown in Figs. 7-9 and 10-11, the culture supernatants of 7D2-E3 and 11C8 increase a proportion of apoptosis cells of Jurkat cells and the culture supernatant of 11C8 increases a proportion of apoptosis cells of HL-60 cells, respectively.

[3] The culture supernatants of 11C8 were used in a coculturing system with HL-60 cells using a feeder layer of cells of the human myeloid stromal cell line KM102, to determine whether these culture supernatants induce apoptosis of HL-60 cells.

Specifically, KM102 cells were seeded in a 2-well Lab-Tek Chamber Slide (Nalge Nunc Intl. Corporation) and brought to a sub-confluent state,  $1 \times 10^5$  cells of HL-60 cells were seeded thereon and cultured for about one day, and then the non-attached HL-60 cells were removed. The

aforementioned culture supernatants were simultaneously added to provide a final concentration of 10% and the cells were cultured for 2 days. After culturing, the cells were fixed with 10% formalin and the apoptosis-induced HL-60 cells were detected by the TUNEL method (ApopTag Plus available from Oncor Inc.). As shown in Figs. 12 and 13, the culture supernatant of 11C8 more increases apoptosis cells of HL-60 cells than the culture supernatant of 9C5 does, which is the culture supernatant of the human IAP non-reacting hybridoma clone used as the control.

#### (5) Antibody purification

For purification of the antibodies produced by hybridomas, the cell lines of the IgG-producing clones 7D2-E3 and 11C8 among the above hybridoma lines were intraperitoneally injected into pristane-administered BALB/c/AnNCrj mice (male, available from Charles River, Japan) according to a standard method. After several weeks, the ascites fluid produced was taken and the antibodies were separated and purified by standard methods. Specifically, the antibodies were purified from the obtained ascites fluid by a Polos Protein A plastic column (Perceptive Biosystems Inc.) and dialyzed with PBS (Dulbecco Inc.), and bands were confirmed with SDS-PAGE analysis. As shown in Fig. 14, electrophoresis using an authentic sample of mouse IgG (Cappel Inc.) as a control confirmed bands for the IgG of clones 7D2-E3 and 11C8 at the same positions as the

authentic sample mouse IgG, under both non-reducing conditions and reducing conditions.

5 In this example, the human Integrin Associated Protein (IAP)-expressing L1210 cells were used as the sensitizing antigen for illustrative purposes, but it is also possible to prepare monoclonal antibodies in the same manner using other human IAP-expressing cells or human IAP itself, and to prepare monoclonal antibodies from an antibody library using the phage display method; this invention is not limited to the aforementioned monoclonal antibodies but encompasses all monoclonal antibodies with properties similar thereto and all hybridomas that produce those monoclonal antibodies.

10 Furthermore, the invention of these monoclonal antibodies also includes humanized antibodies, human antibodies, chimeric antibodies, single-chain antibodies, primatized antibodies and antibody fragments obtained by digesting the antibodies with various enzymes (papain, pepsin, ficin, etc.).

20 The hybridomas producing the monoclonal anti-human Integrin Associated Protein (IAP) antibodies of the invention are novel fused cells created from DBA mice spleen cells and the mouse myeloma cell line P3-U1 as the parent cells; anti-IAP antibody (mouse hybridoma 11C8-F8 (subclone of 11C8), designated as "MABL-1") was deposited as FERM BP-  
25 6100 and anti-IAP antibody (mouse hybridoma 7D2-E3 (subclone of 7D2), designated as "MABL-2") as FERM BP-6101 on

September 1, 1997 with the National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry, located at 1-3 Higashi 1-chome, Tsukuba-shi, Ibaraki-ken, Japan, as an authorized depository for public microorganisms.

Example 2 (Subclass identification of MABL-1 and MABL-2 antibodies)

In order to identify the subclasses of MABL-1 and MABL-2 antibodies obtained above, 500 µl each of MABL-1 and MABL-2 adjusted to 100 ng/ml was spotted on an Isotyping Kit (Stratagene), by which MABL-1 was shown to be IgG1, κ and MABL-2 was shown to be IgG2a, κ.

Example 3 (Human IAP-expressing human leukemia cells)

IAP expression in different human leukemia cell lines was detected by flow cytometry with human IAP-recognizing anti-CD47 antibody (a commercially available product). This antibody was used for the detection because human IAP is believed to be identical to CD47 (Biochem. J., 304, 525-530, 1994). The cell lines used were Jurkat and HL-60 cells (K562 cells, ARH77 cells, Raji cells and CMK cells). The cells were used at  $2 \times 10^5$  cells per sample, the anti-CD47 antibody was incubated with the cells at a final concentration of 5 µg/ml, and the secondary antibody used was FITC-labeled anti-mouse IgG antibody (Becton Dickinson and Company). Mouse IgG1 antibody (Zymed Laboratories Inc.) was used as a control. The results of

the flow cytometry as shown in Fig. 15 (HL-60) and Fig. 16 (Jurkat) confirmed that both cell lines expressed IAP.

Example 4 (Apoptotic effect *in vitro*)

(1) The apoptosis-inducing activity of the MABL-1 and MABL-2 antibodies on L1210 cells transfected with human IAP gene, Jurkat cells and HL-60 cells were examined using Annexin-V (Boehringer Mannheim). The results of analysis with Annexin-V are shown in Figs. 17-22, wherein the dots in the lower left region indicate the live cells, those in the lower right region indicate apoptotic cells, and those in the upper right region indicate necrotic cells. The antibodies used were mouse IgG (Zymed Laboratories Inc.) as a control and MABL-1 and MABL-2 at 10 µg/ml, and after  $4 \times 10^3$  cells of L1210 cells transfected with the human IAP gene were incubated for 72 h and  $6 \times 10^4$  cells of the Jurkat cells were incubated for 48 h, they were analyzed with Annexin-V. Cell death was observed, as shown in Figs. 17-22. For the HL-60 cells, 10 µg/ml of MABL-1 was used, and analysis with Annexin-V at  $1 \times 10^5$  cells likewise revealed cell death.

(2) The apoptosis-inducing activity of MABL-2 antibody Fab fragments on L1210 cells transfected with human IAP gene was examined. Specifically, L1210 cells transfected with human IAP gene were cultured at  $4 \times 10^3$  cells, and MABL-2 antibody Fab fragments and mouse IgG as a control were used at a concentration of 10 µg/ml. The cells were incubated for 72 h and measured with Annexin-V. As a result,

considerable cell death was observed (Figs. 23, 24). The MABL-2 antibody Fab fragments used for the experiment were obtained by digesting the antibody with papain (Pierce Laboratories, Inc.) and purifying it. The MABL-2 antibody Fab fragments were analyzed by SDS electrophoresis (Fig. 25).

Example 5 (Investigation of apoptosis *in vivo*)

(1) Drug efficacy of MABL-1 and MABL-2 (whole IgG)

Human IAP-expressing KPMM2 cells (human myeloma cell line) were transplanted into SCID mice, and on the 10th day after transplantation, MABL-1 and MABL-2 (whole IgG) were administered by single intravenous injection in a dose each of 5 µg/head and 50 µg/head, respectively (n=5); on the 28th day after KPMM2 transplantation, the human IgG levels derived from KPMM2 were measured by ELISA, and the disappearance was confirmed. The survival period was also examined. The results showed marked suppression of blood levels of human IgG in the groups treated with MABL-1 and MABL-2, which represented the anti-tumor effect (Fig. 30). The survival period was also shown to be notably lengthened (Fig. 26).

(2) Drug efficacy of MABL-1 and MABL-2 (F(ab')<sub>2</sub>)

F(ab')<sub>2</sub> fragments prepared by digestion of the MABL-1 and MABL-2 antibodies with pepsin and purification with Protein A (Pierce laboratories, Inc.) were used to examine the anti-tumor effect except the cytotoxic effect via the Fc regions. Specifically, human IAP-expressing

KPMM2 cells (human myeloma cell line) were transplanted into SCID mice, and MABL-1 and MABL-2 F(ab')<sub>2</sub> fragments were intravenously administered to the groups in a dose of 100 µg/head on the 6th and 10th days after transplanting, and to the groups in a dose each of 10 and 30 µg/head on the 6th, 8th and 10th days after transplantation, respectively; the human IgG levels derived from KPMM2 were measured by ELISA on the 30th day after transplantation (Fig. 27). The survival period was also examined up to 90 days after transplanting. As a result, a notable suppressing effect on human IgG levels in the blood was found in the groups treated with MABL-1 and MABL-2, which represented the anti-tumor effect. The survival period was also considerably lengthened (Fig. 28). Fig. 29 shows the SDS electrophoresis pattern for the F(ab')<sub>2</sub> fragments of MABL-1 antibody and MABL-2 antibody.

#### INDUSTRIAL APPLICABILITY

The monoclonal antibodies of this invention are antibodies that specifically recognize human Integrin Associated Protein, and the antigens that induce apoptosis of nucleated blood cells having human Integrin Associated Protein. Accordingly, they are useful as antibodies that recognize human Integrin Associated Protein for its distinction and identification, while also having an action of inducing apoptosis of nucleated blood cells; these properties can be utilized to prepare useful therapeutic

agents in the field of treatment for myeloid leukemia and lymphoid leukemia.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225  
2226  
2227  
2228  
2229  
2230  
2231  
2232  
2233  
2234  
2235  
2